

PATIENT-PERCEIVED EMPATHY FROM NURSES AND THEIR ADHERENCE TO TREATMENT PLANS AMONG PATIENT'S TAKING MAINTENANCE

Maricel S. Quiamco

ABSTRACT

This study determined the significant relationship between patient-perceived empathy from nurses and their adherence to treatment plans using test for correlations and test for differences. The respondents of the study are the 150 patients in the community of Malamba, Marilog District, Davao City taking maintenance medicines who are most willing to participate in the study. Non-probability sampling specifically purposive and quota sampling were sampling technique used. Findings of study revealed that the level of empathy from nurses is very high indicative that majority of the patient's believed that attending nurses are very evidently manifesting empathy towards them. Similarly, the patient's level of adherence to treatment is also very high which implies that patients are complying to their doctor's order regarding their medicine intake. Moreover, test for correlations using Pearson r revealed that no significant relationship was established between the moderating variables (profile age, sex, education, and income), and the patient's overall perceived empathy from nurses (independent variable) from the dependent variable (adherence to treatment plan). However, indicator patient's feedback was found to have a significant positive relationship to patient's adherence to treatment plan. Finally, statistical results using T-test and One Way ANOVA revealed no significant differences were noted in patient's level adherence to treatment plans when analyzed according to their profile.

Keywords: Nurse empathy, adherence, treatment plan, test for correlations, test for differences

INTRODUCTION

Patient-perceived empathy from nurses in fostering trust, reducing anxiety, and promoting adherence to treatment plans. Empathy in healthcare facilitates open communication, patient satisfaction, and holistic care that respects both emotional and physical needs, which ultimately leads to better health outcomes. Patient-centered care highlights the importance of recognizing and respecting patient values, preferences, and experiences, as these factors strengthen the patient-provider relationship and improve compliance with therapeutic plans.

Effective communication between healthcare providers and patients is essential to patient care and recovery, as shown in studies from Malawi and Iran, where patients valued empathetic and respectful communication highly. Poor communication, on the other hand, can negatively affect care quality. Patient adherence remains a significant concern, with approximately 25-50% of patients not following treatment regimens,

affecting health outcomes. Health-care providers, therefore, play a crucial role in educating patients, involving them in decision-making, and supporting adherence to treatment, particularly in chronic diseases like hypertension and diabetes.

In the Philippines, non-adherence is a major cause of treatment failure in chronic diseases, with high non-adherence rates among cardiovascular patients and rural communities facing barriers like financial constraints and low health literacy. This study aims to generate insights to improve patient adherence locally, nationally, and globally.

METHOD

The descriptive correlational design was chosen to examine the relationship between patient-perceived empathy from nurses and treatment adherence. The study took place in Barangay Malamba, Davao City, and included 150 patients (100 on cardiovascular and 50 on diabetes maintenance).

Data was collected using an adapted questionnaire (SPPEN) to measure perceived empathy in three domains: nurse expression, patient feedback, and patient expectations. Adherence was assessed with the Medication Adherence Report Scale (MARS-5), which rates adherence behaviors on a five-point scale. Data gathering involved securing approval, distributing and collecting questionnaires, encoding data, and statistical analysis. Analytical tools included descriptive statistics, Pearson *r* correlation for relationships, and *t*-tests and ANOVA to assess empathy differences across groups.

RESULTS AND DISCUSSION

Demographic Profile

The demographic profile of the 150 respondents reveals that the majority are above 60 years old (44.7%), followed by those aged 51 to 60 years (30%), and the remaining 25.3% are below 50 years old. In terms of sex, the distribution is almost equal, with males comprising 50.7% and females 49.3% of the population. Regarding educational attainment, most respondents reached only the elementary level (68.7%), while 24% completed high school, 4.7% reached college level, and only 2.7% were college graduates. In terms of monthly income, a significant majority (80.7%) earn below Php 10,000, while 18.7% earn between Php 10,001 and Php 20,000, and only 0.7% report an income between Php 20,001 and Php 30,000.

Table 1 Respondent's Demographic Profile

Profile	Frequency	Percent
Age		
Below 50 years	38	25.30
51 – 60 years	45	30.00
Above 60 years	67	44.70
Sex		
Male	76	50.70
Female	74	49.30
Education		
Elementary	103	68.70
High School	36	24.00
College Level	7	4.70
College Graduate	4	2.70
Income		
Below Php10,000	121	80.70
10,001 – 20,000	28	18.70
20,001 – 30,000	1	0.70
Total	150	100.0

Level of Patient's Empathy from Nurses

Table 2 outlines patients' perceptions of empathy from nurses across three indicators: nurse expression, patient feedback, and patient expectations. The overall empathy perception score was very high, with a mean of 4.79 (SD = 0.547), indicating that patients felt strongly that nurses exhibited empathy.

Under nurse expression, patients gave the highest ratings to nurses' attentiveness and friendliness, particularly in statements like "The nurses listen to my opinions" (M = 4.92) and "*The nurses are friendly*" (M = 4.88). However, relatively lower but still very high scores were given to aspects related to emotional understanding, such as "*The nurses can understand my feelings*" (M = 4.24) and "*The nurses care about my daily life*" (M = 4.31). The category mean for this dimension was 4.63, described as very high. This means that nurse expression is always observed.

This result is consistent with earlier studies that emphasize the vital role of empathy in nursing care and its impact on patient satisfaction and communication. For instance, McCabe (2004) found that nurse-patient communication characterized by empathy significantly enhances the quality of care and patient satisfaction. Similarly, Reynolds and Scott (2000) emphasized that empathetic understanding is central to effective therapeutic relationships in nursing, aligning with the current finding that patients believe nurses listen to them, care about their daily life, and understand their emotions. Moreover, Williams and Stickley (2010) found that patients view empathy as a core component of compassionate care, reinforcing the relevance of the current results, where emotional understanding and active listening were rated very high.

In terms of patient's feedback, patients reported a strong willingness to communicate with nurses about their condition ($M = 4.85$) and feelings ($M = 4.57$), as well as a sense of emotional support, as indicated by items like *"The nurses' assistance reduces my depressive symptoms"* ($M = 4.55$). The category mean was 4.66 described as very high. This means suggest that patient's feedback is always observed.

This aligns with existing literature that emphasizes the importance of open communication and emotional support in the nurse-patient relationship. According to O'Hagan et al. (2014), patients who perceive nurses as empathetic and approachable are more likely to share important information about their health and emotional needs, facilitating better clinical outcomes. McCabe (2004) also reported that empathetic communication fosters patient trust and encourages disclosure of sensitive information, contributing to personalized and effective care.

Finally, under patient's expectations, the highest possible ratings were observed, with mean scores of 4.93 for both *"I hope that the nurses taking care of me can patiently listen to me"* and *"...can understand me,"* and 4.91 for *"...can assist me when I am in a bad mood."* This dimension had the category mean of 4.93 described as very high. This denotes that patient's expectations is always observed.

This finding supports the previous studies which emphasizes that patients view empathetic communication and emotional support as fundamental aspects of nursing care. Chan (2002) found that patients' expectations of emotional understanding and patient-centered communication significantly influence their perception of nursing quality. Similarly, Zamanzadeh et al. (2014) reported that patients place high value on nurses' ability to listen attentively, understand their feelings, and provide emotional comfort, especially during moments of vulnerability.

Table 2. Perceived Level of Patients Empathy from Nurses

Patient's Empathy From Nurses	Mean	Std Dev	Description Level
Nurse Expression			
1. I think that the nurses care about me	4.92	0.296	Very High
2. The nurses listen to my opinions	4.85	0.373	Very High
3. The nurses are friendly	4.88	0.326	Very High
4. The nurses aggressively solve my problems	4.55	0.619	Very High
5. The nurses can see things from my perspective	4.31	0.770	Very High
6. The nurses care about my			

daily life	4.24	0.792	Very High
7. The nurses can understand my feelings	4.69	0.586	Very High
Category Level	4.63	0.536	Very High

Patient's Feedback

1. When I am in a bad mood, I will actively tell the nurses	4.85	4.182	Very High
2. I will actively tell the nurses about my condition	4.57	0.548	Very High
3. I am willing to tell the nurses my feelings	4.55	0.574	Very High
4. The nurses' assistance reduces my depressive symptoms	4.45	0.574	Very High
5. I can be encouraged by the nurses taking care of me	4.57	0.523	Very High
Category Level	4.66	1.009	Very High

Patient's Expectations

1. I hope that the nurses taking care of me can patiently listen to me	4.93	0.250	Very High
2. I hope that the nurses taking care of me can understand me	4.93	0.250	Very High
3. When I am in a bad mood, I hope I can obtain assistance from nurses	4.91	0.292	Very High
Category Mean	4.93	0.250	Very High

Overall Mean	4.79	0.547	Very High
---------------------	-------------	--------------	------------------

Level of Patient's Adherence to Treatment Plan

Table 3 shows the level of patient's adherence to treatment plan. The results reveal that the overall mean is 4.14 described as very high. This suggest that patient's adherence to treatment plan is always evident. In particular, the highest mean is 4.46 in

the aspect of *altering the dose*, while the lowest mean is 3.91 in terms of *missing out a dose*.

Table 3. Perceived Level of Patients Adherence to Treatment Plan

Adherence	Mean	Std Dev	Description
1 I take less than instructed	4.40	1.023	Very High
2. I stop taking it for a while	4.09	0.750	Very High
3. I miss out a dose	3.91	0.874	Very High
4. I alter the dose	4.46	0.783	Very High
5. I forget to take it	3.92	2.782	Very High
Overall	4.14	0.941	Very High

Relationship between Perceived Empathy from Nurses and Patient's Adherence to Treatment Plan

The analysis presented in Table 4 shows a test of relationship between empathy and patient's adherence to treatment plan. The overall results reveal that there is no significant relationship between the dimensions of empathy and patient's adherence ($r=.122$, $p>.05$). More specifically, nurse expression and adherence ($r=-.009$, $p > 0.05$), patient expectations and adherence ($r=.154$, $p>.05$).

This finding contrasts with some prior research, which has typically suggested a positive association between perceived empathy and treatment adherence. For instance, Hojat et al. (2010) found that physician empathy was significantly associated with better clinical outcomes and higher adherence, particularly among diabetic patients. Similarly, Zolnieriek and DiMatteo (2009) concluded in their meta-analysis that effective communication, which includes empathy, positively influences patient adherence.

On the other hand, there is a significant relationship between patient feedback and adherence ($r=.185$, $p<.05$). This suggests that when patients actively share information about their condition and feelings with nurses, they are more likely to follow their prescribed treatment regimen. This result is consistent with previous studies that highlight the role of patient engagement and communication in promoting adherence. According to Street et al. (2009), patients who participate actively in conversations with healthcare providers tend to be better informed, more motivated, and more committed to their care plans. Similarly, Haskard Zolnieriek and DiMatteo (2009) found that effective communication especially when patients feel heard and involved significantly improves adherence to treatment.

Table 4. Relationship Between Patient's Empathy from Nurses and Adherence to Treatment Plan

Nurse Empathy	r	p-value	Remarks
Nurse expression	-0.009	0.912	Not Significant
Patient Feedback	0.185	0.024	Significant
Patient Expectations	0.154	0.060	Not Significant
Overall	0.122	0.138	Not Significant

Comparison in the Patients Adherence to Treatment Plan When Analyzed According to their Profile

Using One-Way ANOVA and t-tests, the results showed no significant differences in adherence based on age, gender, education, or income. In terms of age, it revealed no significant difference in adherence across age groups ($F = 2.375$, $p > .05$), also with gender, the t-test showed no significant difference in adherence between males and females ($t = 0.929$, $p > .05$), contrary to Angadi et al. (2020) and Al-Noumani et al. (2022), who found gender-based differences in treatment adherence. Moreover, in terms of education, the results indicated no significant difference in adherence across education levels ($F = 0.871$, $p > .05$), opposing studies that suggest higher education correlates with better adherence (Uchmanowicz, 2018). Furthermore, in the aspect of income, it was found to have no significant difference in adherence across income levels ($F = 1.103$, $p > .05$), contradicting findings by Lee et al. (2019) that associated higher adherence with higher income and employment status.

CONCLUSIONS

1. Respondents are primarily older, with low education and income levels.
2. Patients perceive a high level of empathy from nurses across various indicators.
3. Patients generally adhere well to treatment plans.
4. Only the dimension patient feedback has significant relationship with patient's adherence to treatment plans
5. There is no significant difference in the patient's adherence when grouped according to profile.

REFERENCES

- Abegaz, T. M., Shehab, A., Gebreyohannes, E. A., Bhagavathula, A. S., and Elnour, A. A. (2017). Nonadherence to antihypertensive drugs: A systematic review and meta-analysis. *Med. Baltim.* 92, e5641. doi:10.1097/MD.00000000000005641
- Ágh, T., van Boven, J. F., Wettermark, B., Menditto, E., Pinnock, H., Tsiligianni, I., et al. (2021). A cross-sectional survey on medication management practices for noncommunicable diseases in europe during the second wave of the COVID19 pandemic. *Front. Pharmacol.* 12, 685696. doi:10.3389/fphar.2021.685696
- Al-Noumani H, Al-Harrasi M, Jose J, Al-Naamani Z, Panchatcharam SM. (2022) Medication Adherence and Patients' Characteristics in Chronic Diseases: a National Multi-Center Study. *Clin Nurs Res.* 2022;31(3):426–434.
- Albuquerque A, Howick J. The moral role of clinical empathy in patient healthcare. *Int J Fam Commun Med.* 2023;7(1):11-4.
- Alfian, S. D., Annisa, N., Fajriansyah, F., Perwitasari, D. A., Abdulah, R., Eelko, H., et al. (2020). Modifiable factors associated with non-adherence to antihypertensive or antihyperlipidemic drugs are dissimilar: A multicenter study among patients with diabetes in Indonesia. *J. Gen. Intern. Med.* 1, 2897–2906. –10. doi:10.1007/s11606-020-05809-y
- Alfian, S. D., Denig, P., Coelho, A., and Hak, E. (2019). Pharmacy-based predictors of nonadherence, non-persistence and reinitiation of antihypertensive drugs among patients on oral diabetes drugs in The Netherlands. *PLoS One* 14, e0225390. doi:10.1371/journal.pone.0225390
- Angadi NB, Kavi A, Torgal SS. (2020). Therapeutic Adherence among Elderly with Chronic Diseases Residing in Urban Area of South India–A Descriptive Cross Sectional Study. *Biomed Pharmacol J.* 2020;13(4):2109–2116.
- Archer, J. and Jones, L. (2021), *Theories and Models of Communication*. John Wiley & Sons Ltd. Published 2021 by John Wiley & Sons Ltd.
- Barnlund, Dean C. (5 July 2013). "A Transactional Model of Communication". In Akin, Johnnye; Goldberg, Alvin; Myers, Gail; Stewart, Joseph (eds.). *Language Behavior*. De Gruyter Mouton. pp. 43–61. doi:10.1515/9783110878752.43. ISBN 9783110878752.
- Begum, A., (2023). Meeting Patient Expectations & Ensuring Quality Care As A Nurse. nurses.co.uk

- Berne, E., (2016). Transactional Analysis in Psychotherapy: A Systematic Individual and Social Psychiatry. Pickle Partners Publishing. Retrieved from <https://www.amazon.com/Transactional-Analysis-Psychotherapy-Systemic-Individual/dp/161427844X?asin>
- Chan AHY, Horne R, Hankins M, Chisari C. The Medication Adherence Report Scale: A measurement tool for eliciting patients' reports of nonadherence. *Br J Clin Pharmacol*. 2020;**86**:1281–1288. 10.1111/bcp.14193 [PMC free article] [PubMed] [CrossRef] [Google Scholar]
- Crawford T, Candlin S, Roger P. (2017). New perspectives on understanding cultural diversity in nurse-patient communication. *Collegian*, 2017 Feb 1;24(1):63 – 9. <https://doi.org/10.1016/j.colegn.2015.09.001>
- Creswell, J. W., & Creswell, J. D. (2018). *Research Design: Qualitative, Quantitative, and Mixed Methods Approaches*. Thousand Oaks, CA: Sage Publication, Inc.
- Fourie, M. Using Empathy in Communication. Plus. 2023
- Fuller M, Kamans E, van Vuuren M, Wolfensberger M, de Jong MD. Conceptualizing empathy competence: a professional communication perspective. *Journal of business and technical communication*. 2021 Jul;35(3):333-68
- Gutierrez,M, and Rungpetch Sakulbumrungsil, R. (2023) Effectiveness of a pharmacistled expert system intervention for medication adherence and blood pressure control of adults with hypertension: A randomized controlled trial. <https://doi.org/10.1016/j.sapharm.2023.03.004>Get rights and content.
- Research in Social and Administrative Pharmacy Volume 19, Issue 6, June 2023, Pages 931-943
- Hagerty, T. A., Samuels, W., Norcini-Pala, A., & Gigliotti, E. (2017). Peplau's Theory of Interpersonal Relations: An Alternate Factor Structure for Patient Experience Data?. *Nursing science quarterly*, 30(2), 160–167. <https://doi.org/10.1177/0894318417693286>
- Haribhai-Thompson J, McBride-Henry K, Hales C, Rook H. Understanding of empathetic communication in acute hospital settings: a scoping review. *BMJ open*. 2022 Sep 1;12(9):e06337
- Harris BA, Panozzo G. Therapeutic alliance, relationship building, and communication strategies-for the schizophrenia population: An integrative review. *Arch Psychiatr Nurs*. 2019 Feb;33(1):104-111. doi: 10.1016/j.apnu.2018.08.003. Epub 2018 Aug 8. PMID: 30663612.

- Holmes E.A., Hughes D., Morrison V.L. (2014). Predicting Adherence to Medications Using Health Psychology Theories: A Systematic Review of 20 Years of Empirical Research. *Value Health*. 2014;17:863–876. doi: 10.1016/j.jval.2014.08.2671. [PubMed] [CrossRef] [Google Scholar]
- Johnsson A, Wagman P, Boman A, Pennbrant S. (2018). What are they talking about? Content of the communication exchanges between nurses, patients and relatives in a department of medicine for older people: An ethnographic study. *J Clin Nurs*. 2018 Apr;27(7–9):e1651-9. doi:<https://doi.org/10.1111/jocn.14315>
- Joolae S, Joolaei A, Tschudin V, Bahrani N, Nikbakht NA. Caring relationship: the core component of patients' rights practice as experienced by patients and their companions. *J Med Ethics Hist Med*. 2010;3(4):1–7. [PMC free article] [PubMed] [Google Scholar]
- Kalimashe L, du Plessis E. Mental healthcare users' self-reported medication adherence and their perception of the nursing presence of registered nurses in primary healthcare. *Health SA*. 2021 Jul 22;26:1618. doi: 10.4102/hsag.v26i0.1618. PMID: 34394970; PMCID: PMC8335782
- Kardas P, Dima AL, Potočnjak I, Wettermark B and Agh T (2023), Editorial: Recent advances in attempts to improve medication adherence-from basic research to clinical practice. *Front. Pharmacol*. 14:1144662. doi: 10.3389/fphar.2023.1144662
- Kwame, A., Petrucka, P.M.. (2021). A literature-based study of patient-centered care and communication in nurse-patient interactions: barriers, facilitators, and the way forward. *BMC Nurs* **20**, 158. <https://doi.org/10.1186/s12912-021-00684-2>
- LaMorte, W., (2022). The Health Belief Model. Boston University of Public Health. Retrieved from <https://sphweb.bumc.bu.edu/otlt/MPH-Modules/SB/BehavioralChangeTheories/BehavioralChangeTheories2.html>
- Larsen RE, Pripp AH, Krogstad T, Johannessen Landmark C and Holm LB (2022), Development and validation of a new non-disease-specific survey tool to assess self-reported adherence to medication. *Front. Pharmacol*. 13:981368. doi: 10.3389/fphar.2022.98136
- Lee K.S., Moser D.K., Pelter M.M., Nesbitt T., Dracup K. (2017). Self-care in rural residents with heart failure: What we are missing. *Eur. J. Cardiovasc. Nurs*. 2017;16:326–333. doi: 10.1177/1474515116666439. [PubMed] [CrossRef] [Google Scholar]
- Lin, ChunChih & Chen, LiChin & Han, Chin-Yen & Su, ChingChing & Huang, Ya-Ling. (2022). Exploring the experience of reflective writing among Taiwanese
-

- undergraduate nursing students: A qualitative study. *Journal of Professional Nursing*. 40. 105-110. 10.1016/j.profnurs.2022.03.007.
- Lynch, S., (2022). Adherence to Drug Treatment. Retrieved from <https://www.msdmanuals.com/home/drugs/factors-affecting-response-to-drugs/adherence-to-drug-treatment>
- Madula P, Kalembo WF, Yu H, Kaminga CA. Healthcare provider-patient communication: A qualitative study of women's perceptions during childbirth. *Reprod Health*. 2018 Dec;15(135):1–10. <https://doi.org/10.1186/s12978-018-0580-x>.
- Maffoni M, Traversoni S, Costa E, et al. (2020). Medication adherence in the older adults with chronic multimorbidity: a systematic review of qualitative studies on patient's experience. *Eur Geriatr Med*. 2020;11(3):369–381.
- Magbual, April Monica, (2023) "Effects of Nurse-Patient Relationship on Mental Health Patients' Medication Adherence/Healthcare Outcome". *Nursing | Senior Theses*. 107. <https://doi.org/10.33015/dominican.edu/2023.NURS.ST.28>
- Mamangon, M., Abantao Ds, Bataga Kv, De Ocampo Gg, Domingo Jf, Lazaro Ca, and Tolabing. 2018. "PREVALENCE OF MEDICATION NONADHERENCE AND LEVEL OF HEALTH LITERACY AMONG DIAGNOSED HYPERTENSIVES PATIENTS IN A SELECTED BARANGAY IN MANILA, PHILIPPINES." <https://doi.org/10.6881/ahla.201810.sjp03>.
- Martiningsih W, Winarni S, Acob JR, Baua ME, Nugroho H. Transactional Self Care and Empathy Theory in Nursing (A Perspective). *Open Access Maced J Med Sci [Internet]*. 2021 Nov. 17 [cited 2023 Oct. 18];9(G):273-80. Available from: <https://oamjms.eu/index.php/mjms/article/view/7278>
- McLaren, K., (2021). The Six Essential Aspects of Empathy, Part 4: Perspective Taking. Retrieved from <https://karlamclaren.com/the-six-essential-aspects-of-empathypart-4-perspective-taking/>
- Mokhtar, F., Rajakumar, S., Huri, H., (2023). Adherence tool for prophylactic haemophilia treatment in adult and adolescent patients: A systematic review and metaanalysis protocol. Retrieved from <https://www.researchgate.net/publication/376544017>
- Mondesir F., Levitan, E., Malla, G., Mukerji, R., Carson, A., Safford, M., and Turan, J., (2019). Patient Perspectives on Factors Influencing Medication Adherence Among People with Coronary Heart Disease (CHD) and CHD Risk Factors
- Moore, A., Moore, C., Bunker, L., and Sarnoff, B., (2020). Patient Experience Rounds (PER): Real-time feedback to improve the patient experience and quality of care. *Patient Experience Journal* (The Beryl Institute). Vol. 7, Iss: 3, pp 249253

- Moreno-Poyato, A. R., Montesó-Curto, P., Delgado-Hito, P., Suárez-Pérez, R., AceñaDomínguez, R., Carreras-Salvador, R., ... Roldán-Merino, J. F. (2016). The Therapeutic Relationship in Inpatient Psychiatric Care: A Narrative Review of the Perspective of Nurses and Patients. *Archives of Psychiatric Nursing*, 30(6), 782–787. <https://doi.org/10.1016/j.apnu.2016.03.001>
- Moreno-Poyato AR, Rodríguez-Nogueira Ó; (2021). MiRTCIME.CAT working group. The association between empathy and the nurse-patient therapeutic relationship in mental health units: a cross-sectional study. *J Psychiatr Ment Health Nurs*. 2021 Jun;28(3):335-343. doi: 10.1111/jpm.12675. Epub 2020 Aug 3. PMID: 32657511.
- Moudatsou M, Stavropoulou A, Philalithis A, Koukouli S. The Role of Empathy in Health and Social Care Professionals. *Healthcare (Basel)*. 2020 Jan 30;8(1):26. doi: 10.3390/healthcare8010026. PMID: 32019104; PMCID: PMC7151200.
- Nduaguba SO, Soremekun RO, Olugbake OA, Barner JC. The relationship between patient-related factors and medication adherence among Nigerian patients taking highly active anti-retroviral therapy. *Afr Health Sci*. 2017 Sep;17(3):738745. doi: 10.4314/ahs.v17i3.16. PMID: 29085401; PMCID: PMC5656197.
- Ogle J., Bushnell J.A., Caputi P. Empathy is related to clinical competence in medical care. *Med. Educ*. 2013;47:824–831. doi: 10.1111/medu.12232. [PubMed] [CrossRef] [Google Scholar]
- Orem D. E. (2001). *Nursing concepts of practice* (6th ed.). Mosby
- Peh K.Q.E., Kwan Y.H., Goh H., Ramchandani H., Phang J.K., Lim Z.Y., Loh D.H.F., Østbye T., Blalock D.V., Yoon S., et al. (2021). An Adaptable Framework for Factors Contributing to Medication Adherence: Results from a Systematic Review of 102 Conceptual Frameworks. *J. Gen. Intern. Med*. 2021;36:2784– 2795. doi: 10.1007/s11606-021-06648-1. [PMC free article] [PubMed] [CrossRef] [Google Scholar]
- Roach S. *Caring, the Human Mode of Being: A Blueprint for the Health Professions*. 2nd ed. Ottawa, Ontario: Canadian Hospital Association Press; 2002
- Salzmann-Erikson M., Sjödin M. A narrative meta-synthesis of how people with schizophrenia experience facilitators and barriers in using antipsychotic medication: Implications for healthcare professionals. *Int. J. Nurs. Stud*. 2018;85:7–18. doi: 10.1016/j.ijnurstu.2018.05.003. [PubMed] [CrossRef] [Google Scholar]
- Schöpf AC, Martin GS, Keating MA. Humor as a communication strategy in providerpatient communication in a chronic care setting. *Qual Health Res*. 2017 Feb;27(3):374–90. doi:<https://doi.org/10.1177/1049732315620773>
- Snyder, M., and Haskard-Zolnierrek, K., (2023). *Adherence and Communication. Oxford Bibliographies in Communication*. New York: Oxford Univ. Press.

- Tangkas, K., (2022). Training Standardized Patients to Provide Effective Feedback: Development, Implementation, and its Effect on the Efficacy of Medical Students' Education. PubMed)- Vol. 75, Iss: 10, pp 454-455
- Texas A&M Instrument University, (2019). Why is Empathy Crucial in Nursing?. Retrieved from <https://online.tamui.edu/articles/rnbsn/empathy-is-crucial-in-nursing.aspx>
- Thompson, T., & Haskard-Zolnierrek, K. (2020). Adherence and Communication. *Oxford Research Encyclopedia of Communication*. Retrieved 3 Jun. 2024, from <https://oxfordre.com/communication/view/10.1093/acrefore/9780190228613.001.0001/acrefore-9780190228613-e-23>.
- Tsai, S., Chai, S. and Hwa Wang, H. (2013) Patient-perceived empathy from nurses in Taiwan acute care settings. *Open Journal of Nursing*, **3**, 532-538. doi: 10.4236/ojn.2013.38072.
- Uchmanowicz B, Chudiak A, Uchmanowicz I, Rosińczuk J, Froelicher ES. Factors influencing adherence to treatment in older adults with hypertension. *Clin Interv Aging*. 2018;13:2425.
- Unni E, Bae S. (2022). Exploring a New Theoretical Model to Explain the Behavior of Medication Adherence. *Pharmacy (Basel)*. 2022 Apr 1;10(2):43. doi: 10.3390/pharmacy10020043. PMID: 35448702; PMCID: PMC9025348.
- van der Laan D. M., Elders P. J. M., Boons C. C. L. M., Beckeringh J. J., Nijpels G., Hugtenburg J. G. (2017). Factors associated with antihypertensive medication non-adherence: A systematic review. *J. Hum. Hypertens*. 31, 687–694. 10.1038/jhh.2017.48 [PubMed] [CrossRef] [Google Scholar]
- Waari G, Mutai J, Gikunju J. (2018). Medication adherence and factors associated with poor adherence among type 2 diabetes mellitus patients on follow-up at Kenyatta National Hospital, Kenya. *Pan Af Med J*. 2018;29(1):1–15.
- Zaugg V, Korb-Savoldelli V, Durieux P, Sabatier B. (2018). Providing physicians with feedback on medication adherence for people with chronic diseases taking long-term medication. *Cochrane Database Syst Rev*. 2018 Jan 10;1(1):CD012042. doi: 10.1002/14651858.CD012042.pub2. PMID: 29320600; PMCID: PMC6491069.